

Semisymmetric elementary abelian covers of the Möbius-Kantor graph

Aleksander Malnič,¹ Dragan Marušič^{2,*} Štefko Miklavic² and Primož Potočnik³

University of Ljubljana and Institute of Mathematics, Physics, and Mechanics,
Jadranska 19, 1111 Ljubljana, Slovenia, and
University of Primorska, Cankarjeva 5, 6000 Koper, Slovenia.

Abstract

Let $\varphi_N: \tilde{X} \rightarrow X$ be a regular covering projection of connected graphs with the group of covering transformations isomorphic to N . If N is an elementary abelian p -group, then the projection φ_N is called p -elementary abelian. The projection φ_N is vertex-transitive (edge-transitive) if some vertex-transitive (edge-transitive) subgroup of $\text{Aut} X$ lifts along φ_N , and semisymmetric if it is edge- but not vertex-transitive. The projection φ_N is minimal semisymmetric if φ_N cannot be written as a composition $\varphi_N = \varphi \circ \varphi_M$ of two (nontrivial) regular covering projections, where φ_M is semisymmetric.

Finding elementary abelian covering projections can be grasped combinatorially via a linear representation of automorphisms acting on the first homology group of the graph. The method essentially reduces to finding invariant subspaces of matrix groups over prime fields (see *J. Algebr. Combin.*, **20** (2004), 71–97).

In this paper, all pairwise nonisomorphic minimal semisymmetric elementary abelian regular covering projections of the Möbius-Kantor graph, the Generalized Petersen graph $\text{GP}(8, 3)$, are constructed. No such covers exist for $p = 2$. Otherwise, the number of such covering projections is equal to $(p - 1)/4$ and $1 + (p - 1)/4$ in cases $p \equiv 5, 9, 13, 17, 21 \pmod{24}$ and $p \equiv 1 \pmod{24}$, respectively, and to $(p + 1)/4$ and $1 + (p + 1)/4$ in cases $p \equiv 3, 7, 11, 15, 23 \pmod{24}$ and $p \equiv 19 \pmod{24}$, respectively. For each such covering projection the voltage rules generating the corresponding covers are displayed explicitly.

Keywords: graph, covering projection, lifting automorphisms, homology group, group representation, matrix group, invariant subspaces.

¹University of Ljubljana and Institute of Mathematics, Physics, and Mechanics. Supported in part by “Ministrstvo za visoko šolstvo, znanost in tehnologijo”, research program P1-0285, and by “Ministrstvo za šolstvo, znanost in šport”, research project Z1-3124.

²University of Primorska and Institute of Mathematics, Physics, and Mechanics. Supported in part by “Ministrstvo za visoko šolstvo, znanost in tehnologijo”, research program P1-0285.

³University of Ljubljana and Institute of Mathematics, Physics, and Mechanics. Supported in part by “Ministrstvo za šolstvo, znanost in šport”, research project Z1-4186.

* Corresponding author e-mail: dragan.marusic@guest.arnes.si

1 Introduction

Following the pioneering article of Tutte [28], cubic graphs with specific symmetry properties have been extensively studied over decades by many authors. Much of the work has been focused on classification results, constructions of infinite families, and compiling lists of graphs up to a certain order.

A classification of (connected) cubic *symmetric* (edge- and vertex-transitive) graphs in terms of vertex stabilizers was given by Djoković and Miller [7], and extended to all edge-transitive cubic graphs by Goldschmidt [13]. Foster [2] produced a list of symmetric cubic graphs on up to 512 vertices. Based on Djoković-Miller’s classification, an exhaustive computer search by Conder and Dobcsányi [3] resulted in a complete list of symmetric cubic graphs on up to 768 vertices. Recently, a similar method based on Goldschmidt’s classification was used to compile a list of all cubic *semisymmetric* (edge- but not vertex-transitive) graphs on up to 768 vertices [4].

An important tool in studying symmetry properties of graphs is that of lifting automorphisms along regular covering projections. Recall that a surjective graph morphism $\varphi: \tilde{X} \rightarrow X$ is a *regular N -covering projection* if φ is obtained, roughly speaking, as a quotienting by the action of a semiregular group $N \leq \text{Aut} \tilde{X}$ (see Section 2 for a more precise definition). An automorphism $\alpha \in \text{Aut} X$ *lifts along* $\varphi: \tilde{X} \rightarrow X$ if there exists an automorphism $\tilde{\alpha} \in \text{Aut} \tilde{X}$ such that $\alpha \circ \varphi = \varphi \circ \tilde{\alpha}$. A subgroup $G \leq \text{Aut} X$ *lifts along* φ if each $\alpha \in G$ lifts; the collection of all lifts of all elements of G constitutes a group $\tilde{G} \leq \tilde{X}$, the *lift* of G . A covering projection φ is *vertex-transitive* (*edge-transitive*) if some vertex-transitive (edge-transitive) subgroup of $\text{Aut} X$ lifts along φ , and is *semisymmetric* if it is edge- but not vertex-transitive. Note that if some s -transitive group lifts, then the covering graph is at least s -transitive. It was precisely this observation that led Djoković to construct first examples of infinite families of 5-arc-transitive cubic graphs [6].

The Foster census [2] of symmetric cubic graphs on up to 512 vertices is organized as a “lattice” indicating for a graph, when this information was available, which other graphs in the list are its regular covers. The census of all semisymmetric cubic graphs on up to 768 vertices [4] is organized in a similar fashion; in addition, it also contains information about the maximal groups that lift or project.

The importance of lifting automorphisms along regular coverings was further emphasized in several other recent publications. Regular *elementary abelian* covers, that is, those for which the group of covering transformations is elementary abelian, received particular attention. In [21] it was shown that all cubic graphs admitting a solvable edge-transitive group of automorphisms arise as regular covers of one of the following graphs: the complete graph on four vertices K_4 , the dipole Dip_3 with two vertices and three parallel edges, the complete bipartite graph $K_{3,3}$, the Pappus graph, and the Gray graph (the smallest semisymmetric cubic graph). Moreover, it was shown that each such graph can be obtained from these “basic” graphs by a sequence of edge-transitive elementary abelian regular coverings. A more detailed study of edge- and/or vertex-transitive elementary abelian covers of “small” cubic graphs Dip_3 , K_4 , the cube Q_3 , and $K_{3,3}$ can be found in [9, 10, 11, 22]. Semisymmetric elementary abelian covers of the Heawood graph are considered in [4, 20], while vertex-transitive elementary abelian covers of the Petersen graph can be found in [8, 23].

In this paper we consider edge-transitive and, in particular, semisymmetric elementary abelian regular covering projections of the *Möbius-Kantor graph* $\text{GP}(8,3)$, one of the seven symmetric generalized Petersen graphs [12], the incidence graph of the Möbius-Kantor configuration. The automorphism group of the Möbius-Kantor graph has a rich subgroup structure, which makes the problem considerably more complex than, say, finding covers of the Petersen graph or the Heawood

graph. As with semisymmetric covering projections in general, apart from lifting automorphisms we are here faced with the additional problem of non-lifting automorphisms. Even to find minimal such covering projections (see below for the definition) is a challenging task. We construct all pairwise nonisomorphic minimal semisymmetric elementary abelian regular covering projections of the Möbius-Kantor graph. No such covers exist for $p = 2$. Otherwise, the number of such covering projections is equal to $(p-1)/4$ and $1 + (p-1)/4$ in cases $p \equiv 5, 9, 13, 17, 21 \pmod{24}$ and $p \equiv 1 \pmod{24}$, respectively, and to $(p+1)/4$ and $1 + (p+1)/4$ in cases $p \equiv 3, 7, 11, 15, 23 \pmod{24}$ and $p \equiv 19 \pmod{24}$, respectively. For each such covering projection the voltage rules generating the corresponding covers are displayed explicitly (see Section 7).

For a general theory on lifting automorphisms along elementary abelian covers we refer the reader to [20], where it was shown that the lifting problem essentially reduces to finding invariant subspaces of matrix groups over prime fields, linearly representing the action of automorphisms on the first homology group of the graph. For an alternative approach see [8].




Figure 1: The Möbius-Kantor graph $GP(8,3)$.

2 Preliminaries

Graphs and coverings A *graph* is an ordered pair $X = (V, \sim)$, where $V(X) = V$ is a nonempty set of *vertices* and $\sim$ is an irreflexive symmetric relation on V , called *adjacency*. *Edges* of X are unordered pairs $E(X) = \{uv \mid u \sim v\}$ of adjacent vertices while *arcs* are the corresponding ordered pairs $A(X) = \{(u, v) \mid u \sim v\}$. A *morphism* of graphs $X \rightarrow Y$ is an adjacency-preserving function $V(X) \rightarrow V(Y)$. Composition of morphisms is denoted by $\circ$. In particular, the *automorphism group* $\text{Aut } X$ of X is the subgroup of all adjacency preserving permutations on $V(X)$, equipped with the product $\alpha\beta = \beta \circ \alpha$.

An epimorphism $\wp: \tilde{X} \rightarrow X$ of graphs is called a *regular covering projection* if there exists a subgroup $\text{CT}(\wp) \leq \text{Aut } \tilde{X}$ acting semiregularly (that is, with trivial stabilizers) on $V(\tilde{X})$ such that its orbits coincide with the (*vertex*) *fibres* $\wp^{-1}(v)$, $v \in V(X)$. The *arc fibres* $\wp^{-1}(u, v)$, $u \sim v$,

and the *edge fibres* $\varphi^{-1}(uv)$, $u \sim v$, coincide with the arc and edge orbits of $\text{CT}(\varphi)$. If $\text{CT}(\varphi)$, also known as the group of *covering transformations*, is isomorphic to an abstract group N , then we speak of a regular N -*covering* projection; to emphasize this we sometimes write φ_N instead of just φ . The projection φ_N is *p-elementary abelian* if N is an elementary abelian p -group.

Two regular covering projections $\varphi: \tilde{X} \rightarrow X$ and $\varphi': \tilde{X}' \rightarrow X$ of a graph X are *isomorphic* if there exist an automorphism $\alpha \in \text{Aut } X$ and an isomorphism $\tilde{\alpha}: \tilde{X} \rightarrow \tilde{X}'$ such that $\alpha \circ \varphi = \varphi' \circ \tilde{\alpha}$. In particular, if $\alpha = \text{id}$ then φ and φ' are *equivalent*. If, in the above setting, $\tilde{X} = \tilde{X}'$ and $\varphi = \varphi'$, then we say that α *lifts* (and that $\tilde{\alpha}$ *projects*) along φ . We also say that φ is α -*admissible*.

If G is a subgroup of $\text{Aut } X$ such that $\varphi: \tilde{X} \rightarrow X$ is α -admissible for all $\alpha \in G$, then φ is G -*admissible*. The collection of all such lifts forms a group $\tilde{G} \leq \text{Aut } \tilde{X}$, the *lift* of G . If X and $\tilde{X}$ are both connected, then $\text{CT}(\varphi)$ is precisely the lift of the identity subgroup of $\text{Aut } X$. Moreover, there exists a short exact sequence $\text{CT}(\varphi) \rightarrow \tilde{G} \rightarrow G$. Let φ be G admissible. Then a projection $\varphi': \tilde{X}' \rightarrow X$, isomorphic to φ , need not be G -admissible, although it is admissible for an appropriate subgroup, conjugate to G ; however, any covering projection equivalent to φ is G -admissible. Also, if G^α is conjugate to G , then φ need not be G^α -admissible, although an appropriate projection, isomorphic to φ , is G^α -admissible.

Minimal semisymmetric covers A regular covering projection $\varphi: \tilde{X} \rightarrow X$ is *vertex-transitive* (*edge-transitive*) if some vertex-transitive (edge-transitive) subgroup of $\text{Aut } X$ lifts, and is *semisymmetric* if it is edge- but not vertex-transitive. In order for φ to be vertex-transitive (edge-transitive) it is obviously enough to require that some minimal vertex-transitive (edge-transitive) subgroup lifts. Moreover, if we restrict our considerations up to isomorphism of projections, then, in view of the remarks at the end of the previous subsection, it suffices to consider the above minimal subgroups just up to conjugation.

Let now $\varphi_N: \tilde{X} \rightarrow X$ be a regular N -covering projection. Observe that φ_N is a composition of two regular covering projections if and only if $\varphi = \varphi_{N/K} \circ \varphi_K$, where $K \triangleleft N$. The projection φ is *minimal G -admissible* if φ is G -admissible, and there is no decomposition $\varphi = \varphi' \circ \varphi''$, where φ' and φ'' are regular covering projections such that φ' is G -admissible (see [20, 29] for an extensive discussion). Thus, *minimal vertex-transitive* (*edge-transitive*) regular covering projections are those which cannot be decomposed through “smaller” vertex-transitive (edge-transitive) projections. Similarly, *minimal semisymmetric* covering projections are those semisymmetric ones which cannot be decomposed through “smaller” semisymmetric projections. Note that if a non-vertex-transitive projection is minimal edge-transitive, then it is minimal semisymmetric; however, a minimal semisymmetric projection need not be minimal edge-transitive.

Covers and liftings, combinatorially Let X be a connected graph and N an (abstract) finite group, called the *voltage group*. Assign to each arc of X a *voltage* $\zeta(u, v) \in N$ such that $\zeta(v, u) = (\zeta(u, v))^{-1}$. Let $\text{Cov}(X; \zeta)$ be the *derived graph* with vertex set $V \times N$ and adjacency relation defined by $(u, a) \sim (v, a\zeta(u, v))$, where $u \sim v$ in X . Then the projection onto the first coordinate is a regular N -covering projection $\varphi_\zeta: \text{Cov}(X; \zeta) \rightarrow X$, where the group N , viewed as $\text{CT}(\varphi_\zeta)$, acts via left multiplication on itself. Moreover, it can be shown that each regular N -covering projection $\varphi: \tilde{X} \rightarrow X$ is equivalent to $\varphi_\zeta: \text{Cov}(X; \zeta) \rightarrow X$ for some voltage assignment $\zeta: X \rightarrow N$. Furthermore, if also $\tilde{X}$ is connected, then one can assume that ζ is trivial on the arcs of an arbitrary spanning tree T and that the values on the arcs not in T generate the group N . For an extensive treatment of graph coverings see [14].

The necessary and sufficient condition for φ to be G -admissible can be stated combinatorially in terms of voltages in order for G to lift along an equivalent projection $\text{Cov}(X, \zeta) \rightarrow X$. In particular, if N is an elementary abelian p -group, the following holds [20]. First choose a spanning

tree T of X and a set of arcs $\{x_1, \dots, x_r\} \subseteq A(X)$ containing exactly one arc from each edge in $E(X \setminus T)$. Let $\mathcal{B}_T$ be the corresponding basis of $H_1(X; \mathbb{Z}_p)$ determined by $\{x_1, \dots, x_r\}$. Further, denote by $G^{\#_h} = \{\alpha^{\#_h} \mid \alpha \in G\} \leq \text{GL}(H_1(X; \mathbb{Z}_p))$ the induced action of G on $H_1(X; \mathbb{Z}_p)$, and let $M_G \leq \mathbb{Z}_p^{r,r}$ be the matrix-representation of $G^{\#_h}$ with respect to the basis $\mathcal{B}_T$. By M_G^t we denote the dual group consisting of all transposes of matrices in M_G .

Theorem 2.1 [20, Proposition 6.3, Corollary 6.5] *Let T be a spanning tree of a connected graph X and let the set $\{x_1, x_2, \dots, x_r\} \subseteq A(X)$ contain exactly one arc from each cotree edge. Let $\zeta: A(X) \rightarrow \mathbb{Z}_p^{d,1}$ be a voltage assignment on X which is trivial on T , and let the matrix $Z \in \mathbb{Z}_p^{d,r}$ with columns*

$$\zeta(x_1), \zeta(x_2), \dots, \zeta(x_r)$$

have rank d . Then the following holds.

- (a) *A group $G \leq \text{Aut } X$ lifts along $\wp_\zeta: \text{Cov}(X; \zeta) \rightarrow X$, where the covering graph is connected, if and only if the columns of Z^t form a basis of a M_G^t -invariant d -dimensional subspace $\mathcal{S}(\zeta) \leq \mathbb{Z}_p^{r,1}$.*
- (b) *If $\zeta': A(X) \rightarrow \mathbb{Z}_p^{d,1}$ is another voltage assignment satisfying (a), then $\wp_{\zeta'}$ is equivalent to $\wp_\zeta$ if and only if $\mathcal{S}(\zeta) = \mathcal{S}(\zeta')$. Moreover, $\wp_{\zeta'}$ is isomorphic to $\wp_\zeta$ if and only if there exists an automorphism $\alpha \in \text{Aut } X$ such that the matrix M_α^t maps $\mathcal{S}(\zeta')$ onto $\mathcal{S}(\zeta)$. $\blacksquare$*

By Theorem 2.1, in order to find, up to isomorphism of projections, all vertex-transitive (edge-transitive) elementary abelian covering projections of X , with the derived graph being connected, one has to compute all invariant subspaces of M_H^t , where H ranges through, up to conjugation, all minimal vertex-transitive (edge-transitive) subgroups $H \leq \text{Aut } X$. Next, to actually reduce the respective covering projections up to isomorphism one has to consider the action of $M_{\text{Aut } X}^t$ on the set of all these subspaces, and take just one representative from the corresponding orbits. This also resolves the question of which automorphisms do not lift. It is helpful to note that in finding the orbit of an M_H^t -invariant subspace it is enough to consider just a left transversal of H within $\text{Aut } X$; this is because elements from the same left coset of H map an M_H^t -invariant subspace in the same way. Further, observe that minimal invariant subspaces relative to minimal vertex-transitive (edge-transitive) subgroups correspond to minimal vertex-transitive (edge-transitive) covering projections.

Finally, to obtain the semisymmetric projections, one has to sort out those M_H^t -invariant subspaces (where H ranges over all, up to conjugation, minimal edge-transitive subgroups) which are not invariant for any M_G^t , where $G \leq \text{Aut } X$ is a minimal vertex-transitive subgroup. However, we again remark that the minimal semisymmetric projections might not arise just from those minimal edge-transitive projections which are not vertex-transitive.

Invariant subspaces of matrix groups

As we have seen above, the problem of finding all elementary abelian regular covering projections of a given connected graph, admissible for a given group of automorphisms, is reduced to finding all invariant subspaces of an associated (finite) matrix group over a prime field.

In this context we recall Masche's theorem which states that if the characteristic $\text{Char } \mathbb{F}$ of the field does not divide the order of the group, then the representation is completely reducible. In this case one essentially needs to find just the minimal common invariant subspaces of the generators of the group in question, for the non-minimal subspaces can be expressed as direct sums of some of the minimal ones. (Still, this may involve knowing all invariant subspaces of the generators,

in view of the fact that a minimal invariant subspace for the whole group need not be minimal for neither of the individual generators – although invariant subspaces of a generator are direct sums of the minimal ones for that generator. Additional information about the relations between generators coming from the presentation of the group is beneficial; this is the point where ad-hock techniques are most helpful.) The remaining cases where $\text{Char } \mathbb{F}$ divides the order of the given group could be, technically, more difficult to analyse. But, in contrast with the inherently infinite general problem, there are only finitely many such exceptional field characteristics. Furthermore, with concrete primes one can use computer algebra-packages like MAGMA [1] or GAP [26] with built-in algorithms for computing invariant subspaces. For theoretical background we refer the reader to the work of Holt and Rees [16], and Neumann and Praeger [25].

Clearly, one must be able to find the invariant subspaces of a single matrix in the first place. We recall the following general facts from linear algebra (see for example [17]). Let $A \in \mathbb{F}^{n,n}$ be an $n \times n$ matrix over a field $\mathbb{F}$, acting as a linear transformation $\underline{x} \mapsto A\underline{x}$ on the column vector space $\mathbb{F}^{n,1}$. Denote by $\kappa_A(x) = f_1(x)^{n_1} f_2(x)^{n_2} \dots f_k(x)^{n_k}$ the characteristic polynomial and by $m_A(x) = f_1(x)^{s_1} f_2(x)^{s_2} \dots f_k(x)^{s_k}$ the minimal polynomial of A where $f_j(x)$, $j = 1, \dots, k$, are pairwise distinct irreducible factors over $\mathbb{F}$. Then $\mathbb{F}^{n,1}$ can be written as a direct sum of the A -invariant subspaces

$$\mathbb{F}^{n,1} = \text{Ker } f_1(A)^{s_1} \oplus \text{Ker } f_2(A)^{s_2} \oplus \dots \oplus \text{Ker } f_k(A)^{s_k}.$$

Moreover, all A -invariant subspaces can be found by first considering the invariant subspaces of $\text{Ker } f_j(A)^{s_j}$, $j = 1, \dots, k$, and then taking direct sums of some of these. In particular, the minimal ones are just the minimal A -invariant subspaces of $\text{Ker } f_j(A)^{s_j}$, $j = 1, \dots, k$. The subspace $\text{Ker } f_j(A)^{s_j}$ has dimension $d_j n_j$, where $d_j = \deg f_j(x)$ is the degree of the polynomial $f_j(x)$. Its minimal A -invariant subspaces are cyclic of the form $\langle v, Av, \dots, A^{d_j-1}v \rangle$, where $v \in \text{Ker } f_j(A)$, and each such defines an increasing sequence of length at most s_j of nested invariant subspaces (at least one is precisely of length s_j). If $n_j > s_j$, then a variety of pairwise disjoint minimal cyclic subspaces exist in $\text{Ker } f_j(A)^{s_j}$, and a unique one if $n_j = s_j$. In particular, if $n_j = s_j = 1$, then $\text{Ker } f_j(A)$ itself is the only A -invariant subspace contained in $\text{Ker } f_j(A)$ and hence minimal. Consequently, if $\kappa_A(x) = m_A(x)$ with all $n_j = s_j = 1$, then $\text{Ker } f_j(A)$, $j = 1, \dots, k$, are the only minimal A -invariant subspaces, and all others are direct sums of these.

Finally, the last important issue that we should have addressed is factorisation of polynomials into irreducible factors. Instead, we refer the reader to [18].

3 Transitive subgroups of $\text{Aut GP}(8, 3)$

As intuitively depicted in Figure 1, we identify the vertex-set of the Möbius-Kantor graph $\text{GP}(8, 3)$ with $V = \{1, 2, \dots, 16\}$ and the edge-set with the union of the *outer edges* $E_1 = \{\{i, 1+(i \bmod 8)\} \mid i \in \{1, \dots, 8\}\}$, the *inner edges* $E_2 = \{\{8+i, 8+((i+3) \bmod 8)\} \mid i \in \{1, \dots, 8\}\}$, and the *spokes* $E_3 = \{\{i, 8+i\} \mid i \in \{1, \dots, 8\}\}$.

The automorphism groups of the Generalized Petersen graphs are known [12, 19]. In particular, $G = \text{Aut GP}(8, 3)$ has size 96. In order to describe all its vertex- or/and edge-transitive subgroups we need the following automorphisms, represented as permutations on the vertex set V (see Figures

2 and 3 for a better insight into their actions):

$$\begin{aligned}
\rho &= (1, 2, 3, 4, 5, 6, 7, 8)(9, 10, 11, 12, 13, 14, 15, 16), \\
\sigma &= (1, 14, 7, 12, 5, 10, 3, 16)(2, 11, 8, 9, 6, 15, 4, 13), \\
\psi &= (1, 14, 5, 10)(2, 9, 6, 13)(3, 12, 7, 16)(4, 15, 8, 11), \\
\tau &= (1, 9)(2, 14)(3, 11)(4, 16)(5, 13)(6, 10)(7, 15)(8, 12), \\
\eta &= (1, 9)(2, 12)(3, 15)(4, 10)(5, 13)(6, 16)(7, 11)(8, 14), \\
\omega &= (2, 8, 9)(3, 16, 14)(4, 13, 6)(7, 12, 10).
\end{aligned}$$

Now computations in MAGMA [1] reveal that there are exactly seventeen proper subgroups of $G = \langle \rho, \omega, \sigma \rangle$ which are either vertex- or/and edge-transitive (the respective group-lattice relative to inclusion is shown schematically in Figure 4), namely:




Figure 2: The automorphism ρ is a step-1 rotation, σ is a step-2 rotation, and ψ is a step-4 rotation.




Figure 3: The automorphism τ and σ are reflections across the dashed lines, and ω is a step-4 rotation (with central vertices fixed).

1. There is a unique minimal edge-transitive subgroup, which has order 24 and is hence not vertex-transitive, and a unique maximal edge- but not vertex-transitive subgroup, which has order 48. These are, respectively,

$$H = \langle \rho^2, \omega \rangle \quad \text{and} \quad M = \langle H, \sigma \rangle.$$

Clearly, H and M are normal in G . Observe that H is isomorphic to the semi-direct product $Q \rtimes \mathbb{Z}_3$, where Q is the quaternion group. To see this, identify $\psi = \omega^{-1} \circ \rho^2 \circ \omega$, ρ^2 and

$\rho^2 \circ \psi$ with elements i, j, k of Q , respectively, and ω with the automorphism of Q cyclically permuting i, j and k .

2. There are ten minimal vertex-transitive subgroups, of order 16, which therefore act regularly on vertices and intransitively on edges. These are:
 - Three subgroups of index 2 in the 2-Sylow subgroup $S = \langle \rho, \psi, \tau \rangle$, namely the groups $G_1 = \langle \rho, \psi \rangle$, $G_2 = \langle \rho, \tau \rangle$, and $G_3 = \langle \sigma, \eta \rangle$. Conjugation by ω and ω^2 gives rise to six more subgroups which are contained in the remaining 2-Sylow subgroups S^ω and S^{ω^2} , respectively.
 - The group $G_0 = \langle \rho^2, \psi, \tau \rangle$, which is contained as index 2 subgroup in all three 2-Sylow subgroups, and is moreover normal in G .
3. There are two vertex- and edge-transitive subgroups, namely $K_1 = \langle H, \tau \rangle$ and $K_2 = \langle H, \eta \rangle$. Both are of index 2 in G , and hence act regularly on the set of arcs of $\text{GP}(8, 3)$.




Figure 4: The lattice of vertex- or/and edge-transitive subgroups of $\text{Aut GP}(8, 3)$.

4 Linear representation of generators

Let T be the spanning tree of $\text{GP}(8, 3)$ containing all the spokes $\{i, 8 + i\}$ and all inner edges $\{8 + i, 8 + ((i + 3) \bmod 8)\}$ except for the edge $\{11, 16\}$. Let x denote the dart $(16, 11)$ and let x_i denote the dart $(i, 1 + (i \bmod 8))$, $i \in \{1, 2, \dots, 8\}$. By abuse of notation we identify x and x_i ,

$i = 1, \dots, 8$ with the standard ordered basis of the first homology group arising from the above edges and the spanning tree T .

Let $\rho^{\#_h}, \sigma^{\#_h}, \psi^{\#_h}, \tau^{\#_h}, \eta^{\#_h}$ and $\omega^{\#_h}$ denote the linear transformations induced by the corresponding automorphisms on the homology group $H_1(\text{GP}(8, 3); \mathbb{Z}_p)$, viewed as a vector space over $\mathbb{Z}_p$. Let R, S, P, T, E and O denote their transposed matrices relative to the above standard basis of $H_1(\text{GP}(8, 3); \mathbb{Z}_p)$. A straightforward computation gives

$$\begin{aligned}
R &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad S = \begin{pmatrix} 0 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & -1 & -1 & -1 & -1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & -1 & -1 & -1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & -1 & -1 & -1 & -1 & 0 & 0 & 0 \end{pmatrix} \\
P &= \begin{pmatrix} 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & -1 & -1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & -1 & -1 & -1 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & -1 & -1 \\ 0 & 1 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & -1 & -1 & -1 & 0 \\ 1 & -1 & -1 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 1 & 1 & 0 & 0 & 0 & 1 & 1 & 1 \end{pmatrix} \quad T = \begin{pmatrix} 0 & -1 & -1 & -1 & -1 & -1 & -1 & -1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & -1 & -1 & 0 & 0 & 0 & -1 & -1 & -1 \\ -1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & -1 & 0 & 0 & 0 & -1 & -1 & -1 & -1 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & -1 & -1 & -1 & -1 \end{pmatrix} \\
E &= \begin{pmatrix} 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & -1 & -1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & -1 & -1 \\ 0 & 0 & -1 & -1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 1 \\ 1 & -1 & -1 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & -1 & -1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 \end{pmatrix} \quad O = \begin{pmatrix} 0 & 0 & -1 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & -1 & -1 \\ 1 & 0 & -1 & 0 & 0 & 0 & 0 & 1 & 0 \\ -1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & -1 & -1 & -1 & -1 & -1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \end{pmatrix}.
\end{aligned}$$

5 Calculation of invariant subspaces

In order to find all edge-transitive elementary abelian covers of $\text{GP}(8, 3)$ we need to compute all M_H^t -invariant subspaces, that is, the common invariant subspaces for the matrices R^2 and O . The respective characteristic and minimal polynomials of R^2 and O are

$$\begin{aligned}
\kappa_{R^2}(x) &= -(x-1)(x^4-1)^2, \quad \kappa_O(x) = -(x-1)(x^2+x+1)^4, \\
m_{R^2}(x) &= x^4-1, \quad m_O(x) = x^3-1.
\end{aligned}$$

Observe that their factorization into irreducible factors over $\mathbb{Z}_p$ depends on the congruence class of the prime p modulo 4 and 3, respectively. Indeed, the minimal polynomials have the following factorization:

$$m_{R^2}(x) = \begin{cases} (x-1)^4 & p=2 \\ (x-1)(x+1)(x+i)(x-i) & p \equiv 1 \pmod{4}, i^2 = -1 \\ (x-1)(x+1)(x^2+1) & p \equiv -1 \pmod{4}, \end{cases}$$

$$m_O(x) = \begin{cases} (x-1)^3 & p=3 \\ (x-1)(x-\xi)(x-\xi^2) & p \equiv 1 \pmod{3}, \xi^2 + \xi + 1 = 0 \\ (x-1)(x^2+x+1) & p \equiv -1 \pmod{3}. \end{cases}$$

We first find the R^2 -invariant subspaces, and then sort out those which are also O -invariant. Therefore, the analysis splits into three cases: $p=2$, $p \equiv 1 \pmod{4}$, and $p \equiv -1 \pmod{4}$.

Case $p=2$. In this case the representation of the group H is not completely reducible. First we need an appropriate Jordan basis for the matrix R^2 . Observe that the respective Jordan form has two elementary Jordan matrices of size 4×4 and one of size 1. Thus, a Jordan basis consists of the vectors $b_1, v_1, v_2, v_3, b_2, u_1, u_2, u_3, b_3$, where $R^2 - I$ maps as follows: $v_3 \mapsto v_2 \mapsto v_1 \mapsto b_1 \mapsto 0$, $u_3 \mapsto u_2 \mapsto u_1 \mapsto b_2 \mapsto 0$, and $b_3 \mapsto 0$. Letting $\mathcal{J}_k = \text{Ker}(R^2 - I)^k$ ($k=1, 2, 3, 4$) we have

$$\begin{aligned} \mathcal{J}_1 &= \langle b_1, b_2, b_3 \rangle, \\ \mathcal{J}_2 &= \mathcal{J}_1 \oplus \langle v_1, u_1 \rangle, \\ \mathcal{J}_3 &= \mathcal{J}_2 \oplus \langle v_2, u_2 \rangle, \\ \mathcal{J}_4 &= \mathcal{J}_3 \oplus \langle v_3, u_3 \rangle = \mathbb{Z}_2^{9,1}, \end{aligned}$$

and by computation we find an explicit Jordan basis, for instance:

$$\begin{aligned} b_1 &= (0, 1, 0, 1, 0, 1, 0, 1, 0)^t & b_2 &= (0, 0, 1, 0, 1, 0, 1, 0, 1)^t & b_3 &= (1, 0, 1, 0, 0, 1, 0, 0, 1)^t \\ v_1 &= (0, 1, 0, 0, 0, 1, 0, 0, 0)^t & u_1 &= (0, 0, 1, 0, 0, 0, 1, 0, 0)^t \\ v_2 &= (0, 1, 0, 0, 0, 0, 1, 0, 0)^t & u_2 &= (0, 0, 1, 0, 0, 0, 0, 1, 0)^t \\ v_3 &= (0, 1, 0, 0, 0, 0, 0, 0, 0)^t & u_3 &= (0, 0, 1, 0, 0, 0, 0, 0, 0)^t. \end{aligned}$$

There are many R^2 -invariant subspaces (one can check by MAGMA that there are 322 in all). In particular, the spaces $\mathcal{J}_k$ ($k=1, 2, 3, 4$) are R^2 -invariant. Moreover, $\mathcal{J}_2$, $\mathcal{J}_3$, and of course $\mathcal{J}_4 = \mathbb{Z}_2^{9,1}$, are also $\langle R^2, O \rangle$ -invariant, while $\mathcal{J}_1$ is not. This can easily be checked by acting on the Jordan basis by O . To find all $\langle R^2, O \rangle$ -invariant subspaces, first recall that any O -invariant subspace is a direct sum of minimal O -invariant ones, by Masche's theorem. Since $m_O(x) = (x-1)(x^2+x+1)$, the minimal O -invariant subspaces are: the 1-dimensional eigenspace generated by

$$b = (1, 0, 0, 1, 0, 1, 1, 1, 1) \in \mathcal{J}_2,$$

and the 2-dimensional subspaces $\langle v, Ov \rangle$ where $v \in \text{Ker}(O^2 + O + I)$. We are now going to find the $\langle R^2, O \rangle$ -invariant subspaces in each dimension separately. This is summarized in the table below.

It is easy to see that there are no 1-dimensional $\langle R^2, O \rangle$ -invariant subspaces since $b \notin \mathcal{J}_1$. By computation we have $Ob_1 = b_2$, $Ob_2 = b_1 + b_2$. Thus, $W_2 = \langle b_1, b_2 \rangle$ is $\langle R^2, O \rangle$ -invariant, and hence minimal. Moreover, since $W_2 \setminus \{0\} = (R^2 - I)^{k-1}(\mathcal{J}_k \setminus \mathcal{J}_{k-1})$ ($2 \leq k \leq 4$), every $\langle R^2, O \rangle$ -invariant

subspace W has nontrivial intersection with W_2 . As $W \cap W_2$ cannot be 1-dimensional we have $W_2 \leq W$. Thus, W_2 is the unique minimal $\langle R^2, O \rangle$ -invariant subspace. In particular, W_2 is the unique 2-dimensional one.

Let W be a 3-dimensional $\langle R^2, O \rangle$ -invariant subspace. In view of the above comments on O -invariant subspaces and $W_2 \leq W$ we have that there is a unique 3-dimensional such subspace, namely, $W_3 = W_2 \oplus \langle b \rangle$.

Let W be a 4-dimensional $\langle R^2, O \rangle$ -invariant subspace. Similarly as above, $W = W_2 \oplus \langle w, Ow \rangle$, where $w \in \text{Ker}(O^2 + O + I)$. By taking $w = v_1$ we see, by computation, that $W_4 = W_2 \oplus \langle v_1, Ov_1 \rangle$ is indeed $\langle R^2, O \rangle$ -invariant. We now show that this is the unique such subspace. To this end, consider the intersection $W \cap \mathcal{J}_2$. This intersection is not equal to W_2 because of the action of $R^2 - I$. If $\dim(W \cap \mathcal{J}_2) = 3$, then $W \cap \mathcal{J}_2 = W_3$. But then $b \in W$, a contradiction with $W \leq \text{Ker}(O^2 + O + I)$. Hence $W \leq \mathcal{J}_2$, and so $W \leq \mathcal{J}_2 \cap \text{Ker}(O^2 + O + I)$. Since $b \in \mathcal{J}_2 \setminus \text{Ker}(O^2 + O + I)$, we have that $W = \mathcal{J}_2 \cap \text{Ker}(O^2 + O + I)$, as required.

Let W be a 5-dimensional $\langle R^2, O \rangle$ -invariant subspace. By the above comments on O -invariant subspaces we find that $W = W_3 \oplus \langle w, Ow \rangle$, where $w \in \text{Ker}(O^2 + O + I)$. Note that $W \cap \mathcal{J}_2$ is either $\mathcal{J}_2$ (in which case $W = W_5 = \mathcal{J}_2$), or else this intersection is W_3 . In this latter case $R^2 - I$ takes $\langle w, Ow \rangle$ to $\mathcal{J}_2$ and hence to $W_3 \setminus W_2$. Hence $(R^2 - I)(w) \in b + W_2$ and $(R^2 - I)(Ow) \in b + W_2$ and therefore the difference $(R^2 - I)(w + Ow) \in W_2$. So $w + Ow \in \mathcal{J}_2$. But then also $w = O(w + Ow) \in \mathcal{J}_2$, a contradiction.

Let W be a 6-dimensional $\langle R^2, O \rangle$ -invariant subspace. Similarly as above, $W = W_2 \oplus \langle u, Ou \rangle \oplus \langle w, Ow \rangle$, where $u, w \in \text{Ker}(O^2 + O + I)$. By taking $u = v_1$ and $w = b_3 + v_2$ we find, by computation, that $W_6 = W_4 \oplus \langle b_3 + v_2, Ob_3 + Ov_2 \rangle$ is indeed $\langle R^2, O \rangle$ -invariant. We now show that W must actually be equal to the intersection $\mathcal{J}_3 \cap \text{Ker}(O^2 + O + I)$, which implies that W_6 is the unique such subspace. Clearly, $W \leq \text{Ker}(O^2 + O + I)$. Suppose that $w \notin \mathcal{J}_3$. Then since $\mathcal{J}_3$ is O -invariant, also $Ow \notin \mathcal{J}_3$. Moreover $Ow \neq w$. Consider $(R^2 - I)w$ and $(R^2 - I)(Ow)$. These two vectors are in $\mathcal{J}_3$, and are distinct. Indeed, if $(R^2 - I)(w) = (R^2 - I)(Ow)$, then $R^2(w + Ow) = w + Ow$, implying that $w + Ow \in \mathcal{J}_1$ and hence $w + Ow \in W_2$. But then also $w = O(w + Ow) \in W_2$, a contradiction. Consequently, $W \leq \mathcal{J}_3 \cap \text{Ker}(O^2 + O + I)$. Since $b \notin \text{Ker}(O^2 + O + I)$, we have that the dimension of this intersection is at most 6, forcing $W = \mathcal{J}_3 \cap \text{Ker}(O^2 + O + I)$, as required.

Let W be a 7-dimensional $\langle R^2, O \rangle$ -invariant subspace. Then $W \cap \mathcal{J}_3$ is either $\mathcal{J}_3$ (in which case $W = W_7 = \mathcal{J}_3$), or $W \cap \mathcal{J}_3 = W_5 = \mathcal{J}_2$. In this latter case, take $w \in W \setminus \mathcal{J}_3$. Since $(R^2 - I)w \in \mathcal{J}_3 \setminus \mathcal{J}_2$ we have an immediate contradiction.

Finally, there is no 8-dimensional $\langle R^2, O \rangle$ -invariant subspace, as this could only be $\text{Ker}(O^2 + O + I)$, which, however, is not R^2 -invariant, as one can easily check.

Subspace	Subspace basis
W_2	b_1, b_2

The unique minimal $\langle R^2, O \rangle$ -invariant subspace, $p = 2$.

Subspace	Subspace basis
W_3	b_1, b_2, b
W_4	b_1, b_2, v_1, u_1
W_5	b_1, b_2, v_1, u_1, b
W_6	$b_1, b_2, v_1, u_1, b_3 + v_2, b_3 + u_2$
W_7	$b_1, b_2, v_1, u_1, v_2, u_2, b$

The proper non-trivial and non-minimal $\langle R^2, O \rangle$ -invariant subspaces, $p = 2$.

Case $p \equiv 1 \pmod{4}$. The representation of the group H is completely reducible, by Masche's theorem. Any $\langle R^2, O \rangle$ -invariant subspace is a direct sum of minimal $\langle R^2, O \rangle$ -invariant ones. In turn, each such is a direct sum of the minimal $\langle R^2 \rangle$ -invariant subspaces. So we begin with finding these.

The matrix R^2 is diagonalizable, having the diagonal form $\text{diag}_{R^2}(1, 1, 1, i, i, -1, -1, -i, -i)$. The minimal $\langle R^2 \rangle$ -invariant subspaces are therefore all 1-dimensional subspaces in each of the eigenspaces $\mathcal{L}_{R^2}(1)$, $\mathcal{L}_{R^2}(i)$, $\mathcal{L}_{R^2}(-1)$ and $\mathcal{L}_{R^2}(-i)$. Clearly, the splitting of the whole vector space $\mathbb{Z}_p^{9,1}$ into a direct sum of these minimal subspaces is far from being unique. Denote by $\{v_1, v_2, v_3\}$ and $\{u_1, u_2\}$ the respective ordered bases of $\mathcal{L}_{R^2}(1)$ and $\mathcal{L}_{R^2}(-1)$. Similarly, denote by $\{b_1, b_2\}$ and $\{b_3, b_4\}$ the respective ordered bases of $\mathcal{L}_{R^2}(i)$ and $\mathcal{L}_{R^2}(-i)$. By computation we obtain

$$\begin{aligned} v_1 &= (-2, 1, -1, 1, 1, -1, 1, 1, -1)^t & u_1 &= (0, 1, 0, -1, 0, 1, 0, -1, 0)^t \\ v_2 &= (-4, 1, -3, 1, 1, -3, 1, 1, -3)^t & u_2 &= (0, 0, 1, 0, -1, 0, 1, 0, -1)^t \\ v_3 &= (0, 1, -1, 1, -1, 1, -1, 1, -1)^t \\ b_1 &= (0, 1, 0, i, 0, -1, 0, -i, 0)^t & b_3 &= (0, i, -i-1, 1, i-1, -i, i+1, -1, -i+1)^t \\ b_2 &= (0, 0, 1, 0, i, 0, -1, 0, -i)^t & b_4 &= (0, i+1, -1, -i+1, i, -i-1, 1, i-1, -i)^t. \end{aligned}$$

To find the $\langle R^2, O \rangle$ -invariant subspaces we now look at how O maps the minimal $\langle R^2 \rangle$ -invariant ones.

	v_1	v_2	v_3	u_1	u_2
O	$-u_1$	$-\frac{1}{2}v_2 + \frac{3}{2}v_3$	$-\frac{1}{2}v_2 - \frac{1}{2}v_3$	$-u_2$	v_1

The action of O on the basis $\{v_1, v_2, v_3, u_1, u_2\}$ of $\mathcal{L}_{R^2}(1) \oplus \mathcal{L}_{R^2}(-1)$.

	b_1	b_2	b_3	b_4
O	$\frac{i-1}{2}b_1 + \frac{i+1}{2}b_3$	$\frac{i-1}{2}b_2 + \frac{i+1}{2}b_4$	$\frac{i-1}{2}b_1 - \frac{i+1}{2}b_3$	$\frac{i-1}{2}b_2 - \frac{i+1}{2}b_4$

The action of O on the basis $\{b_1, b_2, b_3, b_4\}$ of $\mathcal{L}_{R^2}(i) \oplus \mathcal{L}_{R^2}(-i)$.

From the above table which shows the action of O on the chosen base vectors we immediately see that the subspaces $\mathcal{L}_{R^2}(1, -1) = \mathcal{L}_{R^2}(1) \oplus \mathcal{L}(-1)$ and $\mathcal{L}_{R^2}(i, -i) = \mathcal{L}(i) \oplus \mathcal{L}_{R^2}(-i)$ are O -invariant. We are now going to identify all minimal subspaces in each of these separately.

In $\mathcal{L}_{R^2}(1, -1)$, the subspace $\langle v_1, u_1, u_2 \rangle$ is obviously the unique minimal $\langle R^2, O \rangle$ -invariant subspace which contains one of the minimal $\langle R^2 \rangle$ -invariant subspaces $\langle v_1 \rangle$, $\langle u_1 \rangle$, or $\langle u_2 \rangle$. Because of its dimension, it is also a unique 3-dimensional such subspace. Next, $\langle v_2, v_3 \rangle$ is also $\langle R^2, O \rangle$ -invariant, and moreover, a complement of $\langle v_1, u_1, u_2 \rangle$. Hence a minimal subspace must contain a vector from $\langle v_2, v_3 \rangle$. Observe that $\langle v_2, v_3 \rangle \leq \mathcal{L}_{R^2}(1)$ is possibly not minimal. That depends on whether O has eigenvectors contained in this subspace. Now the matrix representation of O in the basis $\{v_2, v_3\}$ is

$$\begin{pmatrix} -1/2 & -1/2 \\ 3/2 & -1/2 \end{pmatrix}.$$

If $p \equiv 1 \pmod{3}$ (in addition to $p \equiv 1 \pmod{4}$), then this matrix has two eigenspaces, namely, $\mathcal{L}_O(-\xi) = \langle v_2 + (1 + 2\xi)v_3 \rangle = \langle v_2 + \sqrt{-3}v_3 \rangle$ and $\mathcal{L}_O(\xi) = \langle v_2 - (1 + 2\xi)v_3 \rangle = \langle v_2 - \sqrt{3}v_3 \rangle$, and none if $p \equiv -1 \pmod{3}$. In the first case these two subspaces are minimal $\langle R^2, O \rangle$ -invariant, while in the second case $\langle v_2, v_3 \rangle$ is the only remaining one. We note that in both cases $\mathcal{L}_{R^2}(1, -1)$ splits as the direct sum of minimal $\langle R^2, O \rangle$ -invariant subspaces in a unique way.

As for $\mathcal{L}_{R^2}(i, -i)$, observe that the restriction of O on this subspace has x^2+x+1 as the minimal polynomial. Therefore, the minimal O -invariant subspaces are of the form $\langle w, Ow \rangle$. Moreover, if such a subspace is also R^2 -invariant it must be 2-dimensional. Namely, the way how O maps the base vectors b_1, b_2, b_3 and b_4 implies that for $x \in \mathcal{L}_{R^r}(i)$ we have $Ox - \frac{i-1}{2}x \in \mathcal{L}_{R^r}(-i)$, while for $x \in \mathcal{L}_{R^r}(-i)$ we have $Ox + \frac{i+1}{2}x \in \mathcal{L}_{R^r}(i)$. Hence Ox and x cannot be colinear if x is $\mathcal{L}_{R^2}(i)$ or $\mathcal{L}_{R^2}(-i)$. Moreover, a minimal $\langle R^2, O \rangle$ -invariant subspace must intersect both $\mathcal{L}_{R^2}(i)$ and $\mathcal{L}_{R^2}(-i)$. In fact, by computation we easily see that it must be of the form

$$U(s) = \langle sb_1 + b_2, sb_3 + b_4 \rangle,$$

where $s \in \mathbb{Z}_p \cup \{\infty\}$ (note that $U(\infty) = \langle b_1, b_3 \rangle$). The splitting of $\mathcal{L}_{R^2}(i, -i)$ (and hence of the whole space $\mathbb{Z}_p^{9,1}$) is therefore not unique.

To summarize, in the case $p \equiv 1 \pmod{4}$ the following $\langle R^2, O \rangle$ -invariant subspaces exist, with the non-minimal subspaces being direct sums of the minimal ones.

Subspace	Subspace basis	Contained in	Condition
U_3	v_1, u_1, u_2	$\mathcal{L}_{R^2}(1, -1)$	
U_1	$v_2 + \sqrt{3}v_3$	$\mathcal{L}_{R^2}(1)$	$p \equiv 1 \pmod{12}$
U'_1	$v_2 - \sqrt{3}v_3$	$\mathcal{L}_{R^2}(1)$	$p \equiv 1 \pmod{12}$
U_2	v_2, v_3	$\mathcal{L}_{R^2}(1)$	$p \equiv 5 \pmod{12}$
$U(s), (s \in \mathbb{Z}_p \cup \{\infty\})$	$s b_1 + b_2, s b_3 + b_4$	$\mathcal{L}_{R^2}(i, -i)$	

The minimal $\langle R^2, O \rangle$ -invariant subspaces, $p \equiv 1 \pmod{4}$.

Subspace	Subspace basis	Condition
U_2	v_2, v_3	$p \equiv 1 \pmod{12}$
$U_1 \oplus U(s), (s \in \mathbb{Z}_p \cup \{\infty\})$	$v_2 + \sqrt{3}v_3, s b_1 + b_2, s b_3 + b_4$	$p \equiv 1 \pmod{12}$
$U'_1 \oplus U(s), (s \in \mathbb{Z}_p \cup \{\infty\})$	$v_2 - \sqrt{3}v_3, s b_1 + b_2, s b_3 + b_4$	$p \equiv 1 \pmod{12}$
$U_3 \oplus U_1$	$v_1, u_1, u_2, v_2 + \sqrt{3}v_3$	$p \equiv 1 \pmod{12}$
$U_3 \oplus U'_1$	$v_1, u_1, u_2, v_2 - \sqrt{3}v_3$	$p \equiv 1 \pmod{12}$
$U_2 \oplus U(s), (s \in \mathbb{Z}_p \cup \{\infty\})$	$v_2, v_3, s b_1 + b_2, s b_3 + b_4$	
$\mathcal{L}_{R^2}(i, -i)$	b_1, b_2, b_3, b_4	
$\mathcal{L}_{R^2}(i, -i) \oplus U_1$	$b_1, b_2, b_3, b_4, v_2 + \sqrt{3}v_3$	$p \equiv 1 \pmod{12}$
$\mathcal{L}_{R^2}(i, -i) \oplus U'_1$	$b_1, b_2, b_3, b_4, v_2 - \sqrt{3}v_3$	$p \equiv 1 \pmod{12}$
$\mathcal{L}_{R^2}(1, -1) = U_2 \oplus U_3$	v_1, v_2, v_3, u_1, u_2	
$U_3 \oplus U(s), (s \in \mathbb{Z}_p \cup \{\infty\})$	$v_1, u_1, u_2, s b_1 + b_2, s b_3 + b_4$	
$U_3 \oplus U(s) \oplus U_1, (s \in \mathbb{Z}_p \cup \{\infty\})$	$v_1, u_1, u_2, s b_1 + b_2, s b_3 + b_4, v_2 + \sqrt{3}v_3$	$p \equiv 1 \pmod{12}$
$U_3 \oplus U(s) \oplus U'_1, (s \in \mathbb{Z}_p \cup \{\infty\})$	$v_1, u_1, u_2, s b_1 + b_2, s b_3 + b_4, v_2 - \sqrt{3}v_3$	$p \equiv 1 \pmod{12}$
$\mathcal{L}_{R^2}(i, -i) \oplus U_2$	$b_1, b_2, b_3, b_4, v_2, v_3$	
$\mathcal{L}_{R^2}(1, -1) \oplus U(s), (s \in \mathbb{Z}_p \cup \{\infty\})$	$v_1, v_2, v_3, u_1, u_2, s b_1 + b_2, s b_3 + b_4$	
$\mathcal{L}_{R^2}(i, -i) \oplus U_3$	$v_1, b_1, b_2, u_1, u_2, b_3, b_4$	
$\mathcal{L}_{R^2}(i, -i) \oplus U_3 \oplus U_1$	$v_1, b_1, b_2, u_1, u_2, b_3, b_4, v_2 + \sqrt{3}v_3$	$p \equiv 1 \pmod{12}$
$\mathcal{L}_{R^2}(i, -i) \oplus U_3 \oplus U'_1$	$v_1, b_1, b_2, u_1, u_2, b_3, b_4, v_2 - \sqrt{3}v_3$	$p \equiv 1 \pmod{12}$

The proper non-trivial and non-minimal $\langle R^2, O \rangle$ -invariant subspaces, $p \equiv 1 \pmod{4}$.

Case $p \equiv -1 \pmod{4}$. The matrix R^2 has eigenvalues 1 and -1 , with the respective eigenspaces $\mathcal{L}_{R^2}(1) = \langle v_1, v_2, v_3 \rangle$ and $\mathcal{L}_{R^2}(-1) = \langle u_1, u_2 \rangle$ as in the previous case. The whole space splits into a direct sum of $\langle R^2, O \rangle$ -invariant subspaces $\mathbb{Z}_p^{9,1} = \mathcal{L}_{R^2}(1, -1) \oplus \text{Ker}(R^4 + I)$.

Clearly, the $\langle R^2 \rangle$ -invariant subspaces contained in $\mathcal{L}_{R^2}(1, -1)$ are the same as before. This holds for the $\langle R^2, O \rangle$ -invariant subspaces of $\mathcal{L}_{R^2}(1, -1)$ as well, except for $p = 3$ since the representation of H , restricted to $\mathcal{L}_{R^2}(1, -1)$, is not completely reducible then. Thus, the minimal $\langle R^2, O \rangle$ -invariant subspaces contained in $\mathcal{L}_{R^2}(1, -1)$ are $\langle v_1, u_1, u_2 \rangle$, and either $\langle v_2, v_3 \rangle \leq \mathcal{L}_{R^2}(1)$ or $\langle v_2 + \sqrt{3}v_3 \rangle, \langle v_2 - \sqrt{3}v_3 \rangle \leq \mathcal{L}_{R^2}(1)$ or $\langle v_2 \rangle \leq \mathcal{L}_{R^2}(1)$, depending on whether $p \equiv -1 \pmod{3}$ or $p \equiv 1 \pmod{3}$ or $p = 3$, respectively.

It remains to consider $W = \text{Ker}(R^4 + I)$. The characteristic and the minimal polynomials of R^2 , restricted to W , are $\kappa_{R^2}(x) = (x^2 + 1)^2$ and $m_{R^2}(x) = x^2 + 1$. As $x^2 + 1$ is irreducible, the minimal R^2 -invariant subspaces of W are 2-dimensional, pairwise disjoint, and of the form $\langle w, R^2 w \rangle$. There are $p^2 + 1$ in all. For convenience we denote

$$U_w = \langle w, R^2 w \rangle, \quad w \in W.$$

In order to sort out those U_w which are also O -invariant we have to find $w \in W$ such that $Ow \in U_w$ and $OR^2 w \in U_w$. Actually, it is enough to consider just the first of these two conditions for the second one is then satisfied automatically. To see this, let $Ow = \alpha w + \beta R^2 w$. Observing that the minimal polynomial of O , restricted to W , is $m_O(x) = x^2 + x + 1$, the above equation gives $-w - Ow = O^2 w = \alpha Ow + \beta OR^2 w$. If $\beta \neq 0$, then $OR^2 w \in U_w$, and we are done. It remains to consider the case when $Ow = \alpha w$. Here we need to use the fact that the group H is isomorphic to $Q \rtimes \mathbb{Z}_3$, implying that the generators R^2 and O satisfy the relation $O^{-1}R^2O = OR^2O^{-1}R^2$. Moreover, since $W = \text{Ker}(R^4 + I)$ we have $R^{-2} = -R^2$, and using $O^2 = -O - I$ we finally obtain the relation

$$R^2OR^2O + OR^2 = -R^2.$$

Now, if $Ow = \alpha w$, then $\alpha R^2OR^2w + OR^2w = -R^2w$. This shows that $U_w \cap U_{OR^2w} \neq \emptyset$, forcing $U_w = U_{OR^2w}$, as required.

Note that even if the problem is reduced to finding $w \in W$ such that $Ow \in U_w$, there is still a lot of quite ugly computation involved – even by representing the restrictions of R^2 and O to W as 4×4 matrices, as we shall in fact do later on. But there is an alternative way. Namley, consider the splitting field $\mathbb{F} = \mathbb{Z}_p(i)$, $i^2 = -1$ of the polynomial $x^2 + 1$, and the given matrices over $\mathbb{Z}_p$ acting on $\mathbb{Z}_p(i)^{9,1}$. Then all computations done in the case $p \equiv 1 \pmod{4}$ are valid, except that $i \notin \mathbb{Z}_p$. The 4-dimensional $\langle R^2, O \rangle$ -invariant subspace $W^* = \text{Ker}(R^4 + I)$ of $\mathbb{Z}_p(i)^{9,1}$ splits as $W^* = \mathcal{L}_{R^2}^*(i) \oplus \mathcal{L}_{R^2}^*(-i)$. We have $\mathcal{L}_{R^2}^*(i) = \langle b_1^*, b_2^* \rangle$ where $b_1^* = (0, 1, 0, i, 0, -1, 0, -i, 0)^t$, $b_2^* = (0, 0, 1, 0, i, 0, -1, 0, -i)^t$, and $\mathcal{L}_{R^2}^*(-i) = \langle b_3^*, b_4^* \rangle$ where $b_3^* = (0, i, -i - 1, 1, i - 1, -i, i + 1, -1, -i + 1)^t$, $b_4^* = (0, i + 1, -1, -i + 1, i, -i - 1, 1, i - 1, -i)^t$. The minimal $\langle R^2, O \rangle$ -invariant subspaces of W^* are of the form $U^*(s^*) = \langle s^* b_1^* + b_2^*, s^* b_3^* + b_4^* \rangle$, $s^* \in \mathbb{F} \cup \{\infty\}$. From these subspaces we are now going to derive the minimal $\langle R^2, O \rangle$ -invariant subspaces of W . The latter are precisely those $U^*(s^*)$ which have a basis over $\mathbb{Z}_p$, that is, not involving i . First, consider the vectors

$$\begin{aligned} c_1 &= (0, 1, 0, 0, 0, -1, 0, 0, 0)^t, \\ c_2 &= (0, 0, 1, 0, 0, 0, -1, 0, 0)^t, \\ c_3 &= (0, 0, 0, 1, 0, 0, 0, -1, 0)^t, \\ c_4 &= (0, 0, 0, 0, 1, 0, 0, 0, -1)^t. \end{aligned}$$

Observe that $b_1^* = c_1 + i c_3$ and $b_2^* = c_2 + i c_4$. From $R^2 b_1^* = i b_1^*$ and $R^2 b_2^* = i b_2^*$ we immediately obtain $R^2 c_1 = -c_3$, $R^2 c_3 = c_1$, $R^2 c_2 = -c_4$ and $R^2 c_4 = c_2$. Hence $\{c_1, c_2, c_3, c_4\}$ is a basis for $W = \text{Ker}(R^4 + I)$ over $\mathbb{Z}_p$ and for $W^* = \text{Ker}(R^4 + I)$ over $\mathbb{F}$. The matrix representations of R^2 and O in the ordered basis c_1, c_2, c_3, c_4 (also denoted by the same symbols), are:

$$R^2 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}, \quad O = \begin{pmatrix} -1 & 0 & 1 & 1 \\ 0 & -1 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & -1 & 0 & 0 \end{pmatrix}.$$

Note that direct checking which of the subspaces $U^*(s^*)$ possess a basis over $\mathbb{Z}_p$ still involves a lot of ugly computation. However, there is a short-cut which consists of the following trick. For an arbitrary $w \in W$ over $\mathbb{Z}_p$ let

$$w_* = w - i R^2 w \in W^*.$$

Then $w_* \in \mathcal{L}_{R^2}^*(i)$. In fact, the mapping $w \mapsto w_*$ establishes a bijective correspondence between the minimal R^2 -invariant subspaces U_w in W and the minimal R^2 -invariant subspaces in $\mathcal{L}_{R^2}^*(i) < W^*$ (and hence with all $U^*(s^*)$). This holds true since $z \in U_w$ if and only if $z_* = \lambda^* w_*$ for some $\lambda^* \in \mathbb{F}$, as the reader can check. Thus, in order to find $w \in W$ such that $Ow \in U_w$ we only need to find $w \in W$ such that

$$(Ow)_* = \lambda^* w_*,$$

for some $\lambda^* = \lambda_1 + \lambda_2 i \in \mathbb{F}$. We may further assume that $w_* = s^* b_1^* + b_2^*$, where $s^* \in \mathbb{F} \cup \infty$. (It is precisely this assumption which makes computations in the extension field considerably easier in comparison with computations in the original field.) If $s^* = \infty$, then $w_* = b_1^* = c_1 + i c_3 = c_1 - i R^2 c_1$ and so $w = c_1$. Taking into account the action of O and R^2 we immediately see that $O c_1 \in \langle c_1, R^2 c_1 \rangle$ leads to a contradiction. Thus, no $\langle R^2, O \rangle$ -invariant subspace arises from $U^*(\infty)$. Next, suppose that $s^* = r + t i \in \mathbb{F}$. From $w_* = (r + t i)(c_1 + i c_3) + (c_2 + i c_4)$ we extract the “real part” of w_* in the form $w = r c_1 + c_2 - t c_3$. Since the condition $(Ow)_* = \lambda^* w_*$ is equivalent to $Ow = \lambda_1 w + \lambda_2 R^2 w$, we obtain, after taking into account the action of O on c_1, c_2, c_3 and c_4 , the following system of equations:

$$\begin{aligned} -r - t &= r \lambda_1 - t \lambda_2 \\ t - 1 &= \lambda_1 \\ 1 &= -t \lambda_1 - r \lambda_2 \\ -r - 1 &= -\lambda_2. \end{aligned}$$

So $\lambda_1 = t - 1$, $\lambda_2 = r + 1$, and the parameters r, t must satisfy the condition $r^2 + t^2 + r - t + 1 = 0$ in $\mathbb{Z}_p$. By taking the substitution $r = (-\alpha + \beta - 1)/2$ and $t = (\alpha + \beta + 1)/2$, the condition becomes $\alpha^2 + \beta^2 = -1$. The minimal $\langle R^2, O \rangle$ -invariant subspaces $U_{\alpha, \beta} < W$ therefore arise from $U^*(s^*) < W^*$, where $s^* = (\alpha + 1)\frac{i-1}{2} + \beta\frac{i+1}{2}$. For convenience we also change the basis $\{w, R^2 w\}$ of $U_{\alpha, \beta}$ to

$$\begin{aligned} w_{\alpha, \beta}^1 &= (\alpha + 1) c_1 - c_2 + \beta c_3 + c_4, \\ w_{\alpha, \beta}^2 &= \beta c_1 + c_2 - (\alpha + 1) c_3 + c_4. \end{aligned}$$

Finally, note that the number of subspaces $U_{\alpha, \beta}$ equals $p + 1$, in view of the fact that the equation $\alpha^2 + \beta^2 = -1$, where $p \equiv -1 \pmod{4}$, has $p + 1$ solutions [27, Chp. 4, Proposition 4.4]. Let us comment on this a little bit. Observe that the 2-dimensional orthogonal group over $\mathbb{Z}_p$

(isomorphic to $(\mathbb{Z}_p \cup \{\infty\}, *)$ where $r_1 * r_2 = (r_1 r_2 - 1)/(r_1 + r_2)$) acts regularly on each of its orbits, and the orbits are precisely all vectors with a given norm. Since $p \equiv -1 \pmod{4}$, at least one vector $(\alpha_0, \beta_0)^t$ with norm -1 exists. Hence all solutions of the equation $\alpha^2 + \beta^2 = -1$ can be expressed as

$$(\alpha, \beta) = \begin{pmatrix} \frac{r^2-1}{r^2+1}, & -\frac{2r}{r^2+1} \\ \frac{2r}{r^2+1}, & \frac{r^2-1}{r^2+1} \end{pmatrix} \begin{pmatrix} \alpha_0 \\ \beta_0 \end{pmatrix} \quad (r \in \mathbb{Z}_p \cup \{\infty\}).$$

To summarize, in the case $p \equiv -1 \pmod{4}$ the following $\langle R^2, O \rangle$ -invariant subspaces exist (the case $p = 3$ is listed separately):

Subspace	Subspace basis	Contained in	Condition
U_3	v_1, u_1, u_2	$\mathcal{L}_{R^2}(1, -1)$	
U_1	$v_2 + \sqrt{3}v_3$	$\mathcal{L}_{R^2}(1)$	$p \equiv -5 \pmod{12}$
U'_1	$v_2 - \sqrt{3}v_3$	$\mathcal{L}_{R^2}(1)$	$p \equiv -5 \pmod{12}$
U_2	v_2, v_3	$\mathcal{L}_{R^2}(1)$	$p \equiv -1 \pmod{12}$
$U_{\alpha,\beta}, (\alpha^2 + \beta^2 = -1)$	$w_{\alpha,\beta}^1, w_{\alpha,\beta}^2$	$\text{Ker}(R^4 + I)$	

The minimal $\langle R^2, O \rangle$ -invariant subspaces, $p \equiv -1 \pmod{4}$, $p \neq 3$.

Subspace	Subspace basis	Condition
U_2	v_2, v_3	$p \equiv -5 \pmod{12}$
$U_1 \oplus U_{\alpha,\beta}, (\alpha^2 + \beta^2 = -1)$	$v_2 + \sqrt{3}v_3, w_{\alpha,\beta}^1, w_{\alpha,\beta}^2$	$p \equiv -5 \pmod{12}$
$U'_1 \oplus U_{\alpha,\beta}, (\alpha^2 + \beta^2 = -1)$	$v_2 - \sqrt{3}v_3, w_{\alpha,\beta}^1, w_{\alpha,\beta}^2$	$p \equiv -5 \pmod{12}$
$U_3 \oplus U_1$	$v_1, u_1, u_2, v_2 + \sqrt{3}v_3$	$p \equiv -5 \pmod{12}$
$U_3 \oplus U'_1$	$v_1, u_1, u_2, v_2 - \sqrt{3}v_3$	$p \equiv -5 \pmod{12}$
$U_2 \oplus U_{\alpha,\beta}, (\alpha^2 + \beta^2 = -1)$	$v_2, v_3, w_{\alpha,\beta}^1, w_{\alpha,\beta}^2$	
$\text{Ker}(R^4 + I)$	c_1, c_2, c_3, c_4	
$\text{Ker}(R^4 + I) \oplus U_1$	$c_1, c_2, c_3, c_4, v_2 + \sqrt{3}v_3$	$p \equiv -5 \pmod{12}$
$\text{Ker}(R^4 + I) \oplus U'_1$	$c_1, c_2, c_3, c_4, v_2 - \sqrt{3}v_3$	$p \equiv -5 \pmod{12}$
$\mathcal{L}_{R^2}(1, -1) = U_2 \oplus U_3$	v_1, v_2, v_3, u_1, u_2	
$U_3 \oplus U_{\alpha,\beta}, (\alpha^2 + \beta^2 = -1)$	$v_1, u_1, u_2, w_{\alpha,\beta}^1, w_{\alpha,\beta}^2$	
$U_3 \oplus U_{\alpha,\beta} \oplus U_1, (\alpha^2 + \beta^2 = -1)$	$v_1, u_1, u_2, w_{\alpha,\beta}^1, w_{\alpha,\beta}^2, v_2 + \sqrt{3}v_3$	$p \equiv -5 \pmod{12}$
$U_3 \oplus U_{\alpha,\beta} \oplus U'_1, (\alpha^2 + \beta^2 = -1)$	$v_1, u_1, u_2, w_{\alpha,\beta}^1, w_{\alpha,\beta}^2, v_2 - \sqrt{3}v_3$	$p \equiv -5 \pmod{12}$
$\text{Ker}(R^4 + I) \oplus U_2$	$c_1, c_2, c_3, c_4, v_2, v_3$	
$\mathcal{L}_{R^2}(1, -1) \oplus U_{\alpha,\beta}, (\alpha^2 + \beta^2 = -1)$	$v_1, v_2, v_3, u_1, u_2, w_{\alpha,\beta}^1, w_{\alpha,\beta}^2$	
$\text{Ker}(R^4 + I) \oplus U_3$	$c_1, c_2, c_3, c_4, v_1, u_1, u_2$	
$\text{Ker}(R^4 + I) \oplus U_3 \oplus U_1$	$c_1, c_2, c_3, c_4, v_1, u_1, u_2, v_2 + \sqrt{3}v_3$	$p \equiv -5 \pmod{12}$
$\text{Ker}(R^4 + I) \oplus U_3 \oplus U'_1$	$c_1, c_2, c_3, c_4, v_1, u_1, u_2, v_2 - \sqrt{3}v_3$	$p \equiv -5 \pmod{12}$

The proper non-trivial and non-minimal $\langle R^2, O \rangle$ -invariant subspaces, $p \equiv -1 \pmod{4}$, $p \neq 3$.

Subspace	Subspace basis	Contained in
U	v_2	$\mathcal{L}_{R^2}(1)$
$U_{\alpha,\beta}, (\alpha^2 + \beta^2 = -1)$	$w_{\alpha,\beta}^1, w_{\alpha,\beta}^2$	$\text{Ker}(R^4 + I)$
U_3	v_1, u_1, u_2	$\mathcal{L}_{R^2}(1, -1)$

The minimal $\langle R^2, O \rangle$ -invariant subspaces, $p = 3$.

Subspace	Subspace basis
U_2	v_2, v_3
$U \oplus U_{\alpha,\beta}, (\alpha^2 + \beta^2 = -1)$	$v_2, w_{\alpha,\beta}^1, w_{\alpha,\beta}^2$
$U_3 \oplus U$	v_1, u_1, u_2, v_2
$U_2 \oplus U_{\alpha,\beta}, (\alpha^2 + \beta^2 = -1)$	$v_2, v_3, w_{\alpha,\beta}^1, w_{\alpha,\beta}^2$
$\text{Ker}(R^4 + I)$	c_1, c_2, c_3, c_4
$\text{Ker}(R^4 + I) \oplus U$	c_1, c_2, c_3, c_4, v_2
$\mathcal{L}_{R^2}(1, -1) = U_2 \oplus U_3$	v_1, v_2, v_3, u_1, u_2
$U_3 \oplus U_{\alpha,\beta}, (\alpha^2 + \beta^2 = -1)$	$v_1, u_1, u_2, w_{\alpha,\beta}^1, w_{\alpha,\beta}^2$
$U_3 \oplus U_{\alpha,\beta} \oplus U, (\alpha^2 + \beta^2 = -1)$	$v_1, u_1, u_2, w_{\alpha,\beta}^1, w_{\alpha,\beta}^2, v_2$
$\text{Ker}(R^4 + I) \oplus U_2$	$c_1, c_2, c_3, c_4, v_2, v_3$
$\mathcal{L}_{R^2}(1, -1) \oplus U_{\alpha,\beta}, (\alpha^2 + \beta^2 = -1)$	$v_1, v_2, v_3, u_1, u_2, w_{\alpha,\beta}^1, w_{\alpha,\beta}^2$
$\text{Ker}(R^4 + I) \oplus U_3$	$c_1, c_2, c_3, c_4, v_1, u_1, u_2$
$\text{Ker}(R^4 + I) \oplus U_3 \oplus U$	$c_1, c_2, c_3, c_4, v_1, u_1, u_2, v_2$

The proper non-trivial and non-minimal $\langle R^2, O \rangle$ -invariant subspaces, $p = 3$.

6 Minimal semisymmetric covers

In order to reduce the covering projections up to isomorphism of projections we first have to map the obtained M_H^t -invariant subspaces by the full automorphism group $\text{Aut GP}(8, 3)$ (more precisely, by the corresponding transposed matrix group), and then pick one subspace from each orbit containing an M_H^t -invariant subspace. But since elements in the same (left) coset of M_H^t map each M_H^t -invariant subspace in the same way, it is enough to consider just a (left) transversal of M_H^t within $\text{AutGP}(8, 3)$. One can check that ρ , τ , and σ cover the remaining three cosets of H , distinct from H , and that η lies in the same coset as ρ . As for sorting out the semisymmetric projections we only need to discard those representatives in each orbit which are invariant either for R or for T . This is because the group $M = \langle H, \sigma \rangle$ is semisymmetric while $\langle H, \rho \rangle$ and $\langle H, \tau \rangle$ are vertex-transitive.

Actually, we shall explicitly compute just the minimal semisymmetric covers, in order to keep the long story short.

Case $p = 2$. In Section 5 we computed the Jordan basis $b_1, v_1, v_2, v_3, b_2, u_1, u_2, u_3, b_3$ of $\mathbb{Z}_p^{9,1}$, and found all M_H^t -invariant subspaces. One can check that R maps the above Jordan basis as follows

	b_1	v_1	v_2	v_3	b_2	u_1	u_2	u_3	b_3
R	b_2	$b_2 + u_1$	$u_1 + u_2$	$u_2 + u_3$	b_1	v_1	v_2	v_3	b_3

It is now routine to check that all the M_H^t -invariant subspaces are also R -invariant. Therefore, no 2-elementary abelian covering projection of the Möbius-Kantor graph is semisymmetric. For instance, the covering graph arising from the minimal subspace spanned by $b_1 = (0, 1, 0, 1, 0, 1, 0, 1, 0)^t$ and $b_2 = (0, 0, 1, 0, 1, 0, 1, 0, 1)^t$ is the 2-arc-transitive graph which can be

found in the Foster census under the code F64; the largest subgroup that lifts is actually the whole automorphism group $\text{AutGP}(8, 3)$.

Case $p \equiv 1 \pmod{4}$. By direct computation we get the action of R , T and S on the basis $\{v_1, v_2, v_3, b_1, b_2, u_1, u_2, b_3, b_4\}$. This is given in the table below.

	v_1	v_2	v_3	b_1	b_2	u_1	u_2	b_3	b_4
R	v_1	v_2	$-v_3$	ib_2	b_1	$-u_2$	u_1	$-b_4$	ib_3
T	v_1	$-v_2$	$-v_3$	$-ib_1 + (i-1)b_2$	$-(i+1)b_1 + ib_2$	$-u_1$	$-u_2$	$ib_3 + (-i+1)b_4$	$(i+1)b_3 - ib_4$
S	v_1	$-v_2$	v_3	$(i-1)b_1 + b_2$	$ib_1 - (i-1)b_2$	u_2	$-u_1$	$(i+1)b_3 - ib_4$	$b_3 - (i+1)b_4$

The action of R , T , and S on the basis $\{v_1, v_2, v_3, b_1, b_2, u_1, u_2, b_3, b_4\}$.

The subspaces $\langle v_2 + \sqrt{3}v_3 \rangle$, $\langle v_2 - \sqrt{3}v_3 \rangle \leq \mathcal{L}_{R^2}(1)$ and $\langle v_2, v_3 \rangle \leq \mathcal{L}_{R^2}(1)$ are T -invariant ($\langle v_2, v_3 \rangle \leq \mathcal{L}_{R^2}(1)$ is also R -invariant), and the subspace $\langle v_1, u_1, u_2 \rangle \leq \mathcal{L}_{R^2}(1, -1)$ is R -invariant (as well as T -invariant). Thus, the projections arising from these subspaces are not semisymmetric.

Let us now check the subspaces $U(s)$, $s \in \mathbb{Z}_p \cup \{\infty\}$. From the table above we get that R maps $U(s)$ to $U(-\frac{i}{s})$, that T maps $U(s)$ to $U(\frac{((i-1)/2)s+i}{s-(i-1)/2})$, and that S maps $U(s)$ to $U(\frac{(i-1)s+i}{s-(i-1)})$. This holds true for $s \in \{0, \infty, i-1, (i-1)/2\}$ as well. Moreover, for a given $s \in \mathbb{Z}_p \cup \{\infty\}$ the action of the group $\langle R, T, S \rangle$ on the orbit of $U(s)$ is isomorphic to the action of $\mathbb{Z}_2 \times \mathbb{Z}_2$ and is given in the table below.

	R	T	S
$U(s)$	$U(-\frac{i}{s})$	$U(\frac{((i-1)/2)s+i}{s-(i-1)/2})$	$U(\frac{(i-1)s+i}{s-(i-1)})$
$U(-\frac{i}{s})$	$U(s)$	$U(\frac{(i-1)s+i}{s-(i-1)})$	$U(\frac{((i-1)/2)s+i}{s-(i-1)/2})$
$U(\frac{((i-1)/2)s+i}{s-(i-1)/2})$	$U(\frac{(i-1)s+i}{s-(i-1)})$	$U(s)$	$U(-\frac{i}{s})$
$U(\frac{(i-1)s+i}{s-(i-1)})$	$U(\frac{((i-1)/2)s+i}{s-(i-1)/2})$	$U(-\frac{i}{s})$	$U(s)$

The action of R , T and S on $U(s)$, $s \in \mathbb{Z}_p \cup \{\infty\}$.

Let us now check the fixed points of the action of R , T and S on these subspaces. The only T -invariant subspaces are $U(-1)$ and $U(i)$, and R and S interchange $U(-1)$ and $U(i)$. Thus, the covering projections arising from $U(-1)$ or $U(i)$ are not semisymmetric. The remaining fixed points depend on the congruence class of p modulo 8. We either have $p \equiv 1 \pmod{8}$ or $p \equiv 5 \pmod{8}$.

If $p \equiv 5 \pmod{8}$, then the action of $\langle R, T, S \rangle$ on the orbit of $U(s)$, $s \neq -1, i$, has no fixed points. Consequently, $(p-1)/4$ -pairwise nonisomorphic semisymmetric covering projections arise from $U(s)$, $s \in \mathbb{Z}_p \cup \{\infty\}$. The largest subgroup that lifts is $H = \langle \rho^2, \omega \rangle$.

Suppose now that $p \equiv 1 \pmod{8}$. Then (and only then) there exists $\nu \in \mathbb{Z}_p$ such that $\nu^2 = -i$. In this case the only R -invariant subspaces are $U(\nu)$, $U(-\nu)$, and T and S interchange $U(\nu)$ and $U(-\nu)$. The respective covering projections are not semisymmetric. Next, the only S -invariant subspaces are $U(-1+\nu-\nu^2)$, $U(-1-\nu-\nu^2)$, and R and T interchange $U(-1+\nu-\nu^2)$ and $U(-1-\nu-\nu^2)$. Hence these two are semisymmetric and isomorphic. The maximal subgroup that lifts is $\langle H, \sigma \rangle = M$. If $s \notin \{-1, i, \nu, -\nu, -1+\nu-\nu^2, -1-\nu-\nu^2\}$, then $(p-5)/4$ pairwise nonisomorphic covering projections exists. They are all semisymmetric and the maximal subgroup that lifts is H . Thus, altogether $(p-1)/4$ pairwise nonisomorphic semisymmetric covering projections exist.

Finally, let us check whether among the non-minimal M_H^t -invariant subspaces there exists a minimal semisymmetric one. If a non-minimal M_H^t -invariant subspace U is a direct sum of

two $\langle H, \alpha \rangle$ -invariant subspaces, where $\alpha \in \{\rho, \tau\}$, then U is clearly not semisymmetric, and is discarded. If U is a direct sum of two M_H^t -invariant subspaces, of which one is semisymmetric and the other is vertex-transitive, then even if U is semisymmetric it is not minimal semisymmetric, and is again discarded. Therefore, we only need to check subspaces U being direct sums of an $\langle H, \rho \rangle$ -invariant (but not τ -invariant) subspace and an $\langle H, \tau \rangle$ -invariant (but not ρ -invariant) subspace. In view of these comments, we obtain the minimal semisymmetric covers arising from $\langle v_2 + \sqrt{3}v_3, U(s) \rangle$ and $\langle v_2 - \sqrt{3}v_3, U(s) \rangle$, where $p \equiv 1 \pmod{24}$ and $s \in \{\nu, -\nu\}$ (recall that $\nu^2 = -i$, and that ν exists if and only if $p \equiv 1 \pmod{8}$, while $\langle v_2 \pm \sqrt{3}v_3 \rangle$ exist if and only if $p \equiv 1 \pmod{12}$). Moreover, the respective covering projections are isomorphic since R swaps $v_2 + \sqrt{3}v_3$ with $v_2 - \sqrt{3}v_3$ and T swaps $U(\nu)$ and $U(-\nu)$. The largest group that lifts is H since the respective subspaces are not S -invariant.

Case $p \equiv -1 \pmod{4}$. Like in the previous case, the subspace $\langle v_1, u_1, u_2 \rangle \leq \mathcal{L}_{R^2}(1, -1)$ is R -invariant, the subspaces $\langle v_2 + \sqrt{3}v_3 \rangle$, $\langle v_2 - \sqrt{3}v_3 \rangle \leq \mathcal{L}_{R^2}(1)$, $p \equiv 1 \pmod{3}$, are T -invariant, and $\langle v_2, v_3 \rangle \leq \mathcal{L}_{R^2}(1)$, $p \equiv -1 \pmod{3}$ and $\langle v_2 \rangle$, $p = 3$, are T -invariant and R -invariant. Thus, the covering projections arising from these subspaces are not semisymmetric.

We now consider the subspaces $U_{\alpha, \beta}$, $\alpha^2 + \beta^2 = -1(p)$, spanned (in the ordered basis $\{c_1, c_2, c_3, c_4\}$ of $W = \text{Ker}(R^4 + I)$) by $w_{\alpha, \beta}^1 = (\alpha + 1, -1, \beta, 1)^t$ and $w_{\alpha, \beta}^2 = (\beta, 1, -(\alpha + 1), 1)^t$. Recall that there is a bijective correspondence between the spaces $U_{\alpha, \beta}$ and $U^*(s^*) \leq W^*$, where $s^* = \frac{1}{2}(-\alpha + \beta - 1) + \frac{1}{2}(\alpha + \beta + 1)i$. Hence the action of R , T and S on the subspaces $U_{\alpha, \beta}$ can be derived from the action of R , T and S on the subspaces $U^*(s^*) \leq W^*$.

For instance, $RU^*(s^*) = U^*(-\frac{i}{s^*})$. Now $-\frac{i}{s^*} = -\frac{1}{2\alpha}(\alpha + \beta + 1) + \frac{1}{2\alpha}(\alpha - \beta + 1)i$, and from

$$\frac{1}{2}(-\alpha' + \beta' - 1) = -\frac{1}{2\alpha}(\alpha + \beta + 1)$$

$$\frac{1}{2}(\alpha' + \beta' + 1) = \frac{1}{2\alpha}(\alpha - \beta + 1)$$

we obtain $\alpha' = \frac{1}{\alpha}$ and $\beta' = -\frac{\beta}{\alpha}$. Therefore, $RU_{\alpha, \beta} = U_{\frac{1}{\alpha}, -\frac{\beta}{\alpha}}$. In a similar fashion we obtain the following table:

	R	T	S
$U_{\alpha, \beta}$	$U_{\frac{1}{\alpha}, -\frac{\beta}{\alpha}}$	$U_{\alpha, -\beta}$	$U_{\frac{1}{\alpha}, \frac{\beta}{\alpha}}$
$U_{\frac{1}{\alpha}, -\frac{\beta}{\alpha}}$	$U_{\alpha, \beta}$	$U_{\frac{1}{\alpha}, \frac{\beta}{\alpha}}$	$U_{\alpha, \beta}$
$U_{\alpha, -\beta}$	$U_{\frac{1}{\alpha}, \frac{\beta}{\alpha}}$	$U_{\alpha, \beta}$	$U_{\frac{1}{\alpha}, -\frac{\beta}{\alpha}}$
$U_{\frac{1}{\alpha}, \frac{\beta}{\alpha}}$	$U_{\alpha, -\beta}$	$U_{\frac{1}{\alpha}, -\frac{\beta}{\alpha}}$	$U_{\alpha, \beta}$

The action of R , T and S on $U_{\alpha, \beta}$, $\alpha^2 + \beta^2 = -1(p)$.

As in the case $p \equiv 1 \pmod{4}$ we have that the group $\langle R, T, S \rangle$ acts on these subspaces as the group $\mathbb{Z}_2 \times \mathbb{Z}_2$. Observe that T acts without fixed points. For if T preserves $U_{\alpha, \beta}$, then $\beta = 0$ and therefore $\alpha^2 = -1$, a contradiction since $p \equiv -1 \pmod{4}$. Next, the fixed points of R depend on the congruence class of p modulo 8. We have either $p \equiv 3 \pmod{8}$ or $p \equiv 7 \pmod{8}$. If $p \equiv 7 \pmod{8}$ then there are no fixed points, in view of the fact that -2 is not a square in $\mathbb{Z}_p$ (see [24]). Consequently, $\frac{p+1}{4}$ pairwise nonisomorphic semisymmetric regular covering projections exist in this case. Suppose that $p \equiv 3 \pmod{8}$. Then R preserves $U_{-1, \sqrt{-2}}$ and $U_{-1, -\sqrt{-2}}$, and T and S interchange these two. Thus, the respective covering projections are isomorphic but not semisymmetric. The only two subspaces which are preserved by S are $U_{1, \sqrt{-2}}$ and $U_{1, -\sqrt{-2}}$, and

R and T exchange these two. The largest group that lifts is $\langle H, \sigma \rangle = M$. So the respective two covering projections are isomorphic and semisymmetric. Altogether, $\frac{p+1}{4}$ pairwise nonisomorphic semisymmetric regular covering projections exist. If $(\alpha, \beta) \neq (\pm 1, \pm\sqrt{-2})$, then the largest group that lifts is H .

Finally, let us consider the non-minimal M_H^t -invariant subspaces. Similarly as in the case $p \equiv 1 \pmod{4}$ we find that only four of them are indeed semisymmetric. These are $\langle v_2 + \sqrt{3}v_3, U_{-1,\beta} \rangle$ and $\langle v_2 - \sqrt{3}v_3, U_{-1,\beta} \rangle$, where $p \equiv 19 \pmod{24}$ and $\beta \in \{\sqrt{-2}, -\sqrt{-2}\}$ (recall that -2 is a square in $\mathbb{Z}_p$ if and only if $p \equiv 3 \pmod{8}$, while 3 is a square if and only if $p \equiv 7 \pmod{12}$). Moreover, these four projections are isomorphic since R swaps $v_2 + \sqrt{3}v_3$ and $v_2 - \sqrt{3}v_3$, and T swaps $U_{-1,\sqrt{-2}}$ and $U_{-1,-\sqrt{-2}}$. The largest group that lifts is H since the respective subspaces are not S -invariant.

7 The list

We end the paper by giving the explicit list of all pairwise nonisomorphic minimal semisymmetric elementary abelian regular covering projections of the Möbius-Kantor graph. If $p \equiv 1 \pmod{4}$, then there are $(p-1)/4$ minimal semisymmetric and minimal edge-transitive covering projections; in the subcase $p \equiv 1 \pmod{24}$ there is an additional minimal semisymmetric but not minimal edge-transitive covering projection. If $p \equiv -1 \pmod{4}$, then there are $(p+1)/4$ minimal semisymmetric and minimal edge-transitive covering projections; in the subcase $p \equiv 19 \pmod{24}$ there is an additional minimal semisymmetric but not minimal edge-transitive covering projection. The voltages in $\mathbb{Z}_p^{2,1}$ are trivial on the spanning tree T (see Section 4) while the voltages $\zeta(x)$, $\zeta(x_i)$, $i = 1, \dots, 8$, of the remaining darts are given by the tables below.

THE VOLTAGES

$\zeta(x)$	$\zeta(x_1)$	$\zeta(x_2)$	$\zeta(x_3)$	$\zeta(x_4)$
$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$	$\begin{pmatrix} s \\ 1 + (1+s)i \end{pmatrix}$	$\begin{pmatrix} 1 \\ -1 - s - si \end{pmatrix}$	$\begin{pmatrix} si \\ 1 + s - i \end{pmatrix}$	$\begin{pmatrix} i \\ -s + (1+s)i \end{pmatrix}$

$\zeta(x_5)$	$\zeta(x_6)$	$\zeta(x_7)$	$\zeta(x_8)$
$\begin{pmatrix} -s \\ -1 - (1+s)i \end{pmatrix}$	$\begin{pmatrix} -1 \\ 1 + s + si \end{pmatrix}$	$\begin{pmatrix} -si \\ -1 - s + i \end{pmatrix}$	$\begin{pmatrix} -i \\ s - (1+s)i \end{pmatrix}$

- $p \equiv 5 \pmod{8}$, $i^2 = -1$, $s \in \mathbb{Z}_p$, $s \neq -1, i$. The projections come in groups of four under the action of $\mathbb{Z}_2 \times \mathbb{Z}_2$ on $\mathbb{Z}_p \cup \{\infty\}$ with orbits $\{s, -\frac{i}{s}, \frac{((i-1)/2)s+i}{s-(i-1)/2}, \frac{(i-1)s+i}{s-(i-1)}\}$. The largest group that lifts is H .
- $p \equiv 1 \pmod{8}$, $i^2 = -1$, $s \in \mathbb{Z}_p$, $s \neq -1, i, \nu, -\nu, 1 + \nu - \nu^2, -1 - \nu - \nu^2$, where $\nu^2 = -i$. The projections come in groups of four under the action of $\mathbb{Z}_2 \times \mathbb{Z}_2$ on $\mathbb{Z}_p \cup \{\infty\}$ with orbits $\{s, -\frac{i}{s}, \frac{((i-1)/2)s+i}{s-(i-1)/2}, \frac{(i-1)s+i}{s-(i-1)}\}$. The largest group that lifts is H .
- $p \equiv 1 \pmod{8}$, $i^2 = -1$, $s \in \mathbb{Z}_p$, $s = 1 + \nu - \nu^2$, where $\nu^2 = -i$. The largest group that lifts is M .

THE VOLTAGES

$\zeta(x)$	$\zeta(x_1)$	$\zeta(x_2)$	$\zeta(x_3)$	$\zeta(x_4)$	$\zeta(x_5)$	$\zeta(x_6)$	$\zeta(x_7)$	$\zeta(x_8)$
$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$	$\begin{pmatrix} \alpha + 1 \\ \beta \end{pmatrix}$	$\begin{pmatrix} -1 \\ 1 \end{pmatrix}$	$\begin{pmatrix} \beta \\ -(\alpha + 1) \end{pmatrix}$	$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$	$\begin{pmatrix} -(\alpha + 1) \\ -\beta \end{pmatrix}$	$\begin{pmatrix} 1 \\ -1 \end{pmatrix}$	$\begin{pmatrix} -\beta \\ \alpha + 1 \end{pmatrix}$	$\begin{pmatrix} -1 \\ -1 \end{pmatrix}$

- $p \equiv 7 \pmod{8}$, $\alpha^2 + \beta^2 = -1 \pmod{p}$. The projections come in groups of four under the action of $\mathbb{Z}_2 \times \mathbb{Z}_2$ on the solutions of $\alpha^2 + \beta^2 = -1$ with orbits $\{(\alpha, \beta), (\frac{1}{\alpha}, -\frac{\beta}{\alpha}), (\alpha, -\beta), (\frac{1}{\alpha}, \frac{\beta}{\alpha})\}$. The largest group that lifts is H .
- $p \equiv 3 \pmod{8}$, $\alpha^2 + \beta^2 = -1 \pmod{p}$, $(\alpha, \beta) \neq (\pm 1, \pm\sqrt{-2})$. (Note: empty if $p = 3$.) The projections come in groups of four under the action of $\mathbb{Z}_2 \times \mathbb{Z}_2$ on the solutions of $\alpha^2 + \beta^2 = -1 \pmod{p}$ with orbits $\{(\alpha, \beta), (\frac{1}{\alpha}, -\frac{\beta}{\alpha}), (\alpha, -\beta), (\frac{1}{\alpha}, \frac{\beta}{\alpha})\}$. The largest group that lifts is H .
- $p \equiv 3 \pmod{8}$, $\alpha^2 + \beta^2 = -1 \pmod{p}$, $(\alpha, \beta) = (1, \sqrt{-2})$. The largest group that lifts is M .

THE VOLTAGES

$\zeta(x)$	$\zeta(x_1)$	$\zeta(x_2)$	$\zeta(x_3)$	$\zeta(x_4)$
$\begin{pmatrix} -4 \\ 0 \\ 0 \end{pmatrix}$	$\begin{pmatrix} 1 + \sqrt{3} \\ \nu \\ 1 + (1 + \nu)i \end{pmatrix}$	$\begin{pmatrix} -3 - \sqrt{3} \\ 1 \\ -1 - \nu - \nu i \end{pmatrix}$	$\begin{pmatrix} 1 + \sqrt{3} \\ \nu i \\ 1 + \nu - i \end{pmatrix}$	$\begin{pmatrix} 1 - \sqrt{3} \\ i \\ -\nu + (1 + \nu)i \end{pmatrix}$

$\zeta(x_5)$	$\zeta(x_6)$	$\zeta(x_7)$	$\zeta(x_8)$
$\begin{pmatrix} -3 + \sqrt{3} \\ -\nu \\ -1 - (1 + \nu)i \end{pmatrix}$	$\begin{pmatrix} 1 - \sqrt{3} \\ -1 \\ 1 + \nu + \nu i \end{pmatrix}$	$\begin{pmatrix} 1 + \sqrt{3} \\ -\nu i \\ -1 - \nu + i \end{pmatrix}$	$\begin{pmatrix} -3 - \sqrt{3} \\ -i \\ \nu - (1 + \nu)i \end{pmatrix}$

- $p \equiv 1 \pmod{24}$, $i^2 = -1$, $\nu \in \mathbb{Z}_p$, $\nu^2 = -i$. The largest group that lifts is H .

THE VOLTAGES

$\zeta(x)$	$\zeta(x_1)$	$\zeta(x_2)$	$\zeta(x_3)$	$\zeta(x_4)$	$\zeta(x_5)$	$\zeta(x_6)$	$\zeta(x_7)$	$\zeta(x_8)$
$\begin{pmatrix} -4 \\ 0 \\ 0 \end{pmatrix}$	$\begin{pmatrix} 1 + \sqrt{3} \\ 0 \\ \sqrt{-2} \end{pmatrix}$	$\begin{pmatrix} -3 - \sqrt{3} \\ -1 \\ 1 \end{pmatrix}$	$\begin{pmatrix} 1 + \sqrt{3} \\ \sqrt{-2} \\ 0 \end{pmatrix}$	$\begin{pmatrix} 1 - \sqrt{3} \\ 1 \\ 1 \end{pmatrix}$	$\begin{pmatrix} -3 + \sqrt{3} \\ 0 \\ -\sqrt{-2} \end{pmatrix}$	$\begin{pmatrix} 1 - \sqrt{3} \\ 1 \\ -1 \end{pmatrix}$	$\begin{pmatrix} 1 + \sqrt{3} \\ -1 \\ 0 \end{pmatrix}$	$\begin{pmatrix} -3 - \sqrt{3} \\ -1 \\ -1 \end{pmatrix}$

- $p \equiv 19 \pmod{24}$. The largest group that lifts is H .

We remark that the respective covering graphs need not be themselves semisymmetric. For $p = 3$ we indeed get the unique semisymmetric graph on 144 vertices, and for $p = 5$ we get the unique semisymmetric graph on 400 vertices, see [4]. The case $p = 7$ is still unclear.

References

- [1] W. Bosma, C. Cannon, and C. Playoust, The MAGMA algebra system I: The user language, *J. Symbolic Comput.* **24** (1997), 235–265.
- [2] I. Z. Bouwer (ed.), “The Foster Census”, Charles Babbage Research Centre, Winnipeg, 1988.
- [3] M.D.E. Conder and P. Dobcsányi, Trivalent symmetric graphs on up to 768 vertices, *J. Combin. Math. Combin. Comput.* **40** (2002), 41–63.
- [4] M. D. E. Conder, A. Malnič, D. Marušič, and P. Potočnik, A census of cubic semisymmetric graphs on up to 768 vertices, submitted.
- [5] M. D. E. Conder, A. Malnič, D. Marušič, T. Pisanski, and P. Potočnik, The edge- but not vertex-transitive cubic graph on 112 vertices, *J. Graph Theory*, in print.
- [6] D. Ž. Djoković, Automorphisms of graphs and coverings, *J. Combin. Theory Ser. B* **16** (1974), 243–247.

- [7] D.Ž. Djoković and G.L. Miller, Regular Groups of Automorphisms of Cubic Graphs, *J. Combin. Theory Ser. B* **29** (1980), 195–230.
- [8] S. F. Du, J. H. Kwak, and M. Y. Xu, Lifting of automorphisms on the elementary abelian regular coverings, *Lin. Alg. Appl.*, 373 (2003), 101–119.
- [9] S. F. Du, J. H. Kwak, and M. Y. Xu, On 2-arc transitive covers of complete graphs with covering transformation group $\mathbb{Z}_p^3$, *J. Combin. Theory Ser. B*, to appear.
- [10] Y. Q. Feng and J. H. Kwak, One-regular cubic graphs of order a small number times a prime or a prime square, *J. Aust. Math. Soc.* **76** (2004), 345–356.
- [11] Y. Q. Feng and J. H. Kwak, s -regular cubic graphs as coverings of the complete bipartite graph $K_{3,3}$, *J. Graph Theory* **45** (2004), 101–112.
- [12] R. Frucht, J. E. Graver, and M. E. Watkins, The groups of the generalized Petersen graphs, *Proc. Cambridge Philos. Soc.* **70** (1971), 211–218.
- [13] D. Goldschmidt, Automorphisms of trivalent graphs, *Ann. Math.* **111** (1980), 377–406.
- [14] J. L. Gross and T. W. Tucker, “Topological Graph Theory”, Wiley–Interscience, New York, 1987.
- [15] M. Hofmeister, Graph covering projections arising from finite vector spaces over finite fields, *Discrete Math.* **143** (1995), 87–97.
- [16] D. F. Holt and S. Rees, Testing modules for irreducibility, *J. Austral. Math. Soc. Ser. A* **57** (1994), 1–16.
- [17] N. Jacobson, “Lectures in Abstract Algebra, II. Linear Algebra”, Springer-Verlag, New York, 1953.
- [18] R. Lidl and H. Niederreiter, *Finite Fields*, Encyclopedia of Mathematics and Applications (Cambridge Univ. Press, Cambridge, 1984).
- [19] M. Lovrečič-Saražin, A note on generalized Petersen graphs that are also Cayley graphs, *J. Combin. Theory Ser. B* **69** (1997), 226–229.
- [20] A. Malnič, D. Marušič, and P. Potočnik, Elementary abelian covers of graphs, *J. Algebraic Combin.*, **20** (2004), 71–97.
- [21] A. Malnič, D. Marušič, and P. Potočnik, On cubic graphs admitting an edge-transitive solvable group, *J. Algebraic Combin.*, **20** (2004), 99–113.
- [22] A. Malnič, D. Marušič, P. Potočnik, and C. Q. Wang, An infinite family of cubic edge- but not vertex-transitive graphs, *Discrete Math.* **280** (2004), 133–148.
- [23] A. Malnič and P. Potočnik, Invariant subspaces, duality, and covers of the Petersen graph, *European J. Combin.*, in print.
- [24] T. Nagell, “Euler’s Criterion and Legendre’s Symbol” (in Introduction to Number Theory, pp. 144), Wiley, New York, 1951.
- [25] P. M. Neumann and C. E. Praeger, Cyclic matrices and the Meatax, in *Groups and computation, III (Columbus, OH, 1999)*, 291–300, Ohio State Univ. Math. Res. Inst. Publ., 8, de Gruyter, Berlin, 2001.
- [26] M. Schönert et al., GAP – Groups, Algorithms, and Programming, *Lehrstuhl D für Mathematik*, RWTH, Aachen, 1994.
- [27] C. Small, “Arithmetic of Finite Fields”, Marcel Dekker Inc., New York, 1991.
- [28] W. T. Tutte, A family of cubical graphs, *Proc. Cambridge Phil. Soc.*, **43**(1948), 459–474.
- [29] A. Venkatesh, Covers in imprimitive symmetric graphs, Honours dissertation, Department of Mathematics, University of West Australia, 1997.